

cited in the European Search
Report of EP 02764286.1
Your Ref.: 56246EP006

BEST AVAILABLE COPY



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 1/30, C12Q 1/68, C12N 9/22	A1	(11) International Publication Number: WO 96/09529 (43) International Publication Date: 28 March 1996 (28.03.96)
(21) International Application Number: PCT/EP94/03217 (22) International Filing Date: 24 September 1994 (24.09.94) (71) Applicant (for all designated States except US): BIO-RAD S.P.A. S.R.L. [IT/IT]; Via Morandi, 66, I-20090 Segrate (IT). (72) Inventor; and (75) Inventor/Applicant (for US only): BARBESTI, Silvia [IT/IT]; Via Conca del Naviglio, 10, I-20123 Milano (IT). (74) Agent: RICCARDI, Sergio; Riccardi & Co., Via M. Melloni, 32, I-20129 Milano (IT).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PROCESS OF PRODUCTION OF A PREPARATION FOR STAINING NUCLEAR DNA, PREPARATION OBTAINED THEREBY AND KIT FOR ITS USE (57) Abstract A process for the production of a preparation for specific staining of nuclear DNA is disclosed, that can be used to determine the DNA cell contents by flow cytometry. The working solution of such a preparation, containing besides the specific dye, a thermolabile enzyme and a surfactant, is dehydrated under vacuum so as to stabilize it and to modify its characteristic of access to ligand and hydration, and in this way it can be stored for a long time, preferably packed in single dose vials, and then reconstituted at the moment of use, maintaining its properties perfectly unaltered.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

"PROCESS OF PRODUCTION OF A PREPARATION FOR STAINING NUCLEAR DNA, PREPARATION OBTAINED THEREBY AND KIT FOR ITS USE"

The present invention relates to a process for the production of a preparation for staining nuclear DNA to be used in flow cytometry, the preparation obtained
5 by such a process, and a kit for its use.

It is well known that cytofluorometric analysis took a fundamental role in diagnostics and its use is steadily increasing, together with the development of new and more sophisticated analysis techniques and apparatus.

More particularly, the determination of cellular DNA contents through flow
10 cytometry is fundamental in the oncologic field as well as in determining the cellular ploidy by cytofluorometric analysis.

Up to now the working solution containing the specific dye and a thermolabile enzyme, was prepared in an extemporaneous way at the time of using it, because of the well known stability problems of the proteins in the
15 anhydrous condition, but such a process is not at all satisfactory because it is also well known that extemporaneous preparations depend too much from the present situation and the operator's skill, thus they do not warrant reproducible results.

Indeed, as proteins in the anhydrous condition generally show stability problems and their rehydration is rather long, difficult and sometimes incomplete,
20 it was hitherto considered impossible to make a preparation for staining nuclear DNA to be stored in the anhydrous condition and to be reconstructed at the time of use.

It was now surprisingly found that this problem is brilliantly solved by mixing the protein with a surfactant, so as to cause the formation of a water envelope
25 around the molecule, obtain a better stability with time and a more complete

hydration at the moment of use.

In other words, not a true lyophilization but instead a crystallization of the preparation is carried out, so that it may be perfectly stored for a long time.

The ingredients to be used in the production of the preparation according to the present invention must be chosen obviously in a proper way. As enzyme a specific ribonuclease is used, for instance type A bovine ribonuclease; as a dye, a fluorescent halide binding specifically to nucleic acids, preferably propidium iodide, but other fluorescent halides are suitable as well; as a surfactant, it is necessary to use a non ionic surfactant such as NONIDET P 40 (BDH Laboratory
10 Supplies Ltd., Poole, Dorset, Great Britain).

Once the working solution is prepared, it is delivered in vials in such amount as to allow an analytic determination for each vial, which is therefore single dose, and the solution water is evaporated under vacuum.

It is known that DNA staining may take place either in a hypotonic medium, if it is desired to stain only isolated nuclei, or in an isotonic medium, if it desired
15 to keep intact the cell structure for possible morphologic studies or checking proliferation markers.

Two different diluents are therefore prepared for reconstituting the working solution, consisting of salts with a different concentration according to either
20 diluent.

Consequently, three different single dose kits may be supplied to users; namely the vial containing the preparation, accompanied by a bottle containing either the isotonic or the hypotonic diluent, or two vials each with one of the diluents.

25 The production of the preparation according to the present invention will be

BEST AVAILABLE COPY

now illustrated in detail in the following practical example of preparation, which however should not be construed as limiting in any way the scope of the invention.

EXAMPLE

- 5 The chosen dye is propidium iodide, that links specifically by intercalation to the nucleic acids in their double strand structure; it is therefore necessary to remove the cell RNA by enzymatic digestion, using a specific ribonuclease, so as to avoid an overestimation (moreover the RNA contents of a cell is not constant, but it may vary with the state of activity).
- 10 Access of the dye to the nuclear compartment is made easier by using a surfactant, present at a low concentration, which makes the nuclear membrane permeable.
- 15 Firstly a high concentration stock solution of propidium iodide is prepared, dissolving 500 $\mu\text{g/ml}$ of the dye in twice distilled water. The mixture so obtained is left standing on a magnetic agitator for at least 4-6 hours so as to obtain the complete dissolution of the dye. The solution is then filtered with a membrane of cellulose acetate with a pore size 0.22 μ so as to remove possible insoluble residues. Subsequently the working solution is prepared: to one part of stock solution, 9 parts of twice distilled water are added (the final concentration of the dye results to be 50 $\mu\text{g/ml}$), 1 mg/ml of crystallized bovine ribonuclease type A and 0.1% v/v NONIDET P 40. The mixture is again placed on a magnetic agitator for at least 2 hours, adjusting the magnet speed so as not to create foam, and then filtered with a membrane of cellulose acetate with a pore size 0.22 μ . The working solution is delivered in conic bottom test tubes having a capacity of 1.5 ml, in a
- 25 quantity of 1 ml for each tube. Water evaporation is obtained using a thickener

with a fixed angle rotor, with a low speed centrifugation (1500 rpm) for 8 hours at controlled temperature (25°C) under vacuum. At the end the test tubes are closed with a proper pressure stopper.

Two different diluents are then prepared for the reconstitution of the working
5 solution. Isotonic diluent No. 1: 0.155 M NaCl, 15 mM NaN_3 in twice distilled water. Hypotonic diluent No. 2: 0.01 M NaCl, 15 mM NaN_3 in twice distilled water. Both solution, after complete dissolution on magnetic agitator, are filtered with a membrane of cellulose acetate with a pore size of 0.22μ for removing possible insoluble residues and sterilization from bacterial contaminants.

BEST AVAILABLE COPY**CLAIMS**

1) Process of production of a preparation for staining nuclear DNA, constituted by a working solution containing a specific dye and a thermolabile enzyme for RNA digestion, characterized by the steps of mixing said enzyme with
5 a surfactant and evaporating water of the working solution under vacuum, so as to obtain a crystallization residue that may be stored for a long time and reconstituted at the moment of use by suitable diluents.

2) Process according to Claim 1, characterized in that a fluorescent halide is used as specific dye.

10 3) Process according to Claim 2, characterized in that propidium iodide is used as fluorescent halide.

4) Process according to Claim 1, characterized in that a non ionic surfactant is used as a surfactant.

5) Process according to Claim 4, characterized in that the product marketed
15 under the name NONIDET P40 is used as non ionic surfactant.

6) Process according Claim 1, characterized in that a specific ribonuclease is used as enzyme.

7) Process according to Claim 6, characterized in that bovine ribonuclease type A is used as ribonuclease.

20 8) Process according to anyone of Claims 1 to 7, characterized in that a hypotonic or isotonic medium is used as a diluent for the reconstitution of the working solution at the moment of use.

9) Preparation for staining nuclear DNA, characterized in that it comprises the crystallization residue of a working solution containing a specific dye, a
25 thermolabile enzyme for RNA digestion and an enzyme stabilizing surfactant, to

be reconstituted at the moment of use by means of a hypotonic or isotonic medium.

10) Preparation for staining nuclear DNA, whenever obtain through the process according to one or more of Claims 1 to 8.

5 11) Preparation for staining nuclear DNA according to Claim 9 or 10, characterized in that it is packed in single dose vials allowing only one analytic determination for each vial.

12) Kit for using the preparation of Claim 11, characterized in that the single dose vial is accompanied by a bottle containing either the isotonic or the hypotonic
10 diluent, or by two bottles each containing one of said two diluents.

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/EP 94/03217

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 G01N1/30 C12Q1/68 C12N9/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 G01N C12Q C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EUROPEAN JOURNAL OF HISTOCHEMISTRY, vol. 37, 1993 page 68 BARBESTI ET AL. 'dna flow cytometry performed by stabilised propidium iodide staining' see the whole document ---	1-7
A A	US,A,4 087 328 (SWAISGOOD) 2 May 1978 see column 7, line 64 - line 66 see column 2, line 50 - column 3, line 12 --- -/--	1 8,9

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

15 June 1995

Date of mailing of the international search report

30.06.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Hocquet, A

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/EP 94/03217

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>IMMUNOPHARMACOLOGY, vol. 28, no. 3, 18 October 1994 AMSTERDAM, pages 241-257, A.BROWN ET AL. 'propellant -driven aerosols of functional proteins' see page 242, column 1, line 46 - column 2, line 38 see page 253, column 2, line 4 - page 254, column 1, line 9</p> <p style="text-align: center;">---</p>	1
A	<p>BIOCHIMICA ET BIOPHYSICA ACTA , vol. 947, 1988 pages 209-246, LUISI ET AL. 'reverse micelles as hosts for proteins and small molecules' see page 232, column 2, last paragraph - page 235, column 2, paragraph 3 see page 241, paragraph XI - page 242</p> <p style="text-align: center;">---</p>	1
A	<p>US,A,4 668 618 (THORNTHWAITE) 26 May 1987 see column 2, line 2 - line 20 see column 4, line 7 - line 49 see column 6, line 39 - line 43</p> <p style="text-align: center;">-----</p>	2-5

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat Application No

PCT/EP 94/03217

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4087328	02-05-78	NONE	
US-A-4668618	26-05-87	US-A- 4906561	06-03-90

THIS PAGE BLANK (USPTO)